



Original Research Article

One Pot Reductive Synthesis of Benzimidazole Derivatives from 2-Nitro Aniline and Aromatic Aldehydes Using Zn/NaHSO₃ in Water Medium

Hridoydip Ranjan Dasgupta, Suvodip Mukherjee, Pranab Ghosh*

Department of Chemistry, University of North Bengal, Darjeeling 734013, India

ARTICLE INFO

Article history

Submitted: 2020-08-25

Revised: 2020-10-03

Accepted: 2020-10-07

Available online: 2020-10-24

Manuscript ID: [PCBR-2008-1126](#)

DOI: [10.22034/pcbr.2020.245413.1126](#)

KEYWORDS

2-nitroaniline,
one pot reaction,
Reductive cyclisation reaction,
Benzimidazole

ABSTRACT

Small amount of Zn dust and NaHSO₃ was utilized to efficiently synthesize benzimidazole derivatives via one pot reductive cyclocondensation process in water medium at 100°C temperature. Very good to excellent yields in reasonably short reaction times, high atom economy and usage of readily available starting material, operational simplicity and easy workup are the fundamental features of this protocol.

HIGHLIGHT

- One pot reaction
- Chemoselective reductive cyclisation reaction.
- Aqueous medium.
- Shorter reaction time.
- Easy reaction set-up.
- Inexpensive, easily available reagent, easily separable by simple filtration.
- Mild reaction condition.


* Corresponding author: Pranab Ghosh

✉ E-mail: pizy12@gmail.com

Tel.: +91 (0353) 2776381; fax: +91 (0353) 2699001.

© 2020 by SPC (Sami Publishing Company)

GRAPHICAL ABSTRACT



Introduction

Substituted benzimidazoles display beneficial biological properties such as antihypertensives, antivirals, antifungals, anticancers, and antihistaminics[1]. Furthermore, compounds belonging to this class have been employed as versatile building blocks of anthelmintics[2], proton pump inhibitors[3], and a host of pharmaceutical compounds [4]. A series of compounds with benzimidazole motifs have been used in the textile industry as wetting, emulsifying, foaming, or softening agents or as dispersants for use in dyeing.

As a consequence, a number of researchers have set their goal to synthesize benzimidazole derivatives. Historically, the first benzimidazole was prepared in 1872 by Hoebrecker. To overcome the drawbacks of classical methods, the most common approaches to benzimidazoles are the combination of 1, 2-phenylenediamine with aldehyde [5-26], carboxylic acid [27-31]

and ester [32-33] under different catalytic conditions.

Most of the protocols, however, suffer from drawbacks such as harsh reaction conditions, high temperature, strongly acidic / basic condition, prolonged reaction time, use of homogeneous catalyst, low yield and suffer from rapid loss of catalytic activity. Although the acceptable yield of benzimidazoles has been reported in most of the protocols where either toxic metal catalyst or costly reagents were used, people had to suffer handling tedious reaction conditions and work up process. To avoid these limitations, it is imperative to develop a high yielding greener, simple, cost effective and catalyst free efficient method for its synthesis with a broad range of substrate applicability.

Although 2-nitroaniline appears to be a potential precursor for a possible route to benzimidazole derivatives but except very few [34-36] reports, people have not yet explored its applicability.





Figure 1: Examples of some bioactive benzimidazole derivatives

Based on the above reports, it seems necessary to develop an easy, cost-efficient and simple method for the preparation of these derivatives which meet the present demand for sustainable development. Keeping these issues in mind, we here reported a simple and mild one-pot method for the synthesis of 2-substituted benzimidazoles (scheme 1, d) from 2-nitroanilines by reductive cyclocondensation with a combination of suitable

metal, Zn and metal salt NaHSO_3 at 100°C in water. NaHSO_3 is a non-hazardous, easily available, inexpensive weakly acidic species having pKa value 6.97, which helps to trigger the reaction in the presence of Zn. Its adduct formation ability with the aldehydes may contribute to cyclocondensation in the present protocol.



Scheme 1: Comparison of previous work and our work for the synthesis of benzimidazole derivatives.

Table 2. ^aReaction (Scheme-2b) conditions optimization.

Entry	Metal	Solvent	Time(min.)	Temperature(°C)	Yield (%) ^b
1	Fe	DMF	60	100	68
2	Fe	DMSO	60	100	60
3	Fe	Toluene	60	100	56
4	Fe	H ₂ O	60	100	80
5	Fe	-	60	100	Nil
6	Cu	H ₂ O	60	100	75
7	Zn	H ₂ O	60	100	94
8	-	H ₂ O	60	100	Nil
9	Zn	H ₂ O	60	RT	Nil
10	Zn	H ₂ O	60	80	50
11	Zn	H ₂ O	60	60	Nil
12	Zn	H ₂ O	60	40	Nil
13	Zn	H ₂ O	90	100	95
14	Zn	H ₂ O	45	100	87
15	Zn	H ₂ O	120	100	94

^aReaction of *o*-nitrobenzaldehyde (1 mmol), Zn (3mmol), NaHSO₃ (6 mmol) in water on magnetic stirrer. ^bIsolated yield of benzimidazole

Table 3. ^aOptimization of amount of Zn and NaHSO₃

Entry	Zn (mmol)	NaHSO ₃ (mmol)	Time (min)	Yield (%) ^b
1	3	3	60	54
2	3	4	60	62
3	3	5	60	82
4	3	6	60	94
5	3	7	60	94
6	2	6	60	75

^aReaction of *o*-nitrobenzaldehyde (1mmol), Zn (2-3 mmol), NaHSO₃ (4-7 mmol) in water on magnetic stirrer at 100°C.

^bIsolated yield.

























To screen the reaction, 2-nitroaniline and benzaldehyde were selected as model substrates for intended transfiguration. Initially, we performed the reaction of *o*-nitroaniline with benzaldehyde in water at room temperature for 1h on a magnetic stirrer in presence of the combination of Fe powder and sodium bisulphite. The reaction yielded only the diamine (*o*-phenylenediamine). The yield of diamine decreases with a rise in temperature and benzimidazole was undetected (Table 1, scheme-2a). It was also tried (scheme-2a) without using NaHSO₃ (Table 1, entry 4), but it failed to produce the diamine even at a trace amount. Thus, it is obvious that, as a hydrogen ion's source, NaHSO₃ plays a vital role to reduce nitro to amine. With this experimental data we followed our scheme 2b. In this process *o*-nitroaniline was reduced to 1, 2-diamine with Zn and NaHSO₃ in presence of water at room temperature and the process was completed within 5 minutes. It was followed by the addition of benzaldehyde with continuous stirring at 100 °C. As a solvent, water was first screened (Scheme 2b, Table 2, entry 4), and very surprisingly no product was isolated in its absence (Table 2, entry 5). Further, as the most easily available and most significant, its green nature has really enriched the objective of the present investigation.

The presence of metal is the necessary requirement for the initial reduction of the nitro compound (Table 2, entry 8) and in comparison to Fe and Cu, Zn is estimable in terms of yield of the product and the time of completion of reaction (Table 2, entries 4, 6, 7). We also tried the scheme at different temperature; finally, at 100 °C temperature the desired product, i.e. benzimidazole, was isolated as a single compound (Table 2, entry 7). Further, we optimized the amount of Zn and NaHSO₃ required (Table 3).


Further investigation towards the optimization of the process revealed that a 3 mmol Zn and 6 mmol NaHSO₃, (Table 3) under atmospheric pressure at 100°C yielded the best result to produce the desired benzimidazole in 1h (compared with Table 2, entries 7, 14, 15). This procedure was followed for all of the reactions listed in Table (4). Staggering part of our reaction was chemoselective reduction of nitro to amine. We really enthralled and exhilarated on observing Table 4 entry 3, 4, 5, 6, 7, 8 that reducible groups remain intact after completion of the reaction. The reaction took place smoothly to produce corresponding benzimidazole in moderate to high yields.

From the above observation, we can propose a possible mechanism and tentative intermediates for the above developed protocol for the synthesis of benzimidazole as follows (scheme-3)

Table 4. ^aZn and NaHSO₃ mediated reduction to amines

Entry	Reactant	Product	Time (min)	Yield (%) ^b	Melting point ^c (°C)
1			50	93	289-290 ^[37]
2			70	83	>290
3			80	90	200-202 ^[37]
4			50	85	238-240 ^[37]
5			45	89	254-255 ^[37]
6			90	87	290-292 ^[38]
7			60	90	277-279 ^[37]
8			70	94	289
9			45	93	196-198 ^[39]
10			80	90	240-242 ^[38]
11			50	87	>290 ^[40]
12			60	89	267-270 ^[40]

^aReaction of nitro compound (1mmol), Zn (3 mmol), NaHSO₃ (6 mmol) in water at 100 °C for different time intervals on magnetic stirrer. ^bIsolated yields. ^cMelting points of the isolated compounds.



Scheme 3: Proposed mechanism for the synthesis of benzimidazole derivatives

Conclusion

We have developed a novel and efficient protocol through one pot reductive cyclocondensation of 2-nitroaniline with aromatic aldehydes to benzimidazole with Zn/NaHSO₃ in water. The fascinating part of our method in comparison to the conventional methods is its simplicity, cost effectiveness, environmentally benign approach and a less time consuming process. We also earn that Zn/NaHSO₃ in water is also a better chemoselective reducing system to reduce nitro to amine. Thus, it could potentially be

complementary to follow the existing methods for the synthesis of biologically active benzimidazoles moiety.

Acknowledgment

One of us (HRD) is thankful to UGC, New Delhi, India for financial support.

References

- [1] G. L. Gravatt, B. C. Baguley, W. R. Wilson, W. A. Denny, DNA-Directed Alkylating Agents. 6. Synthesis and Antitumor Activity of DNA Minor Groove-Targeted Aniline Mustard Analogues of Pibenzimol (Hoechst 33258),

- Journal of Medicinal Chemistry, 37 (1994) 4338-4345.
- [2] R. K. Prichard, S. Ranjan, Anthelmintics, *Veterinary Parasitology*, 46 (1993) 113-120
- [3] J. Horn, The proton-pump inhibitors: similarities and differences, *Clinical Therapeutics*, 22 (2000) 266-280.
- [4] J. Bhaumik, Z. Yao, K. E. Borbas, M. Taniguchi, J. S. Lindsey, Masked Imidazolyl-Dipyrromethanes in the Synthesis of Imidazole-Substituted Porphyrins. *The Journal of Organic Chemistry*, 71 (2006) 8807-8817.
- [5] V. D. Patil, J. Patil, P. Rege, G. Dere, Mild and Efficient Synthesis of Benzimidazole Using Lead Peroxide Under Solvent-Free Conditions, *Synthetic Communication*, 41 (2011) 58-62.
- [6] F. Hakimi, M. Fallah-Mehrjardi, E. Golrasan, Yttrium Aluminum Garnet (YAG: $Al_5Y_3O_{12}$) as an Efficient Catalyst for the Synthesis of Benzimidazole and Benzoxazole Derivatives, *Chemical Methodologies*, 4(2020) 234-244.
- [7] M. Asif, Green Synthesis of Benzimidazole Derivatives: an Overview on Green Chemistry and Its Applications, *Chemical Methodologies*, 3 (2019) 620-631.
- [8] A. Nozarie, Metal-Organic Framework MIL-53(Fe) as a highly efficient reusable catalyst for the synthesis of 2-aryl-1H-benzimidazole, *Chemical Methodologies*, 3 (2019) 704-712.
- [9] S. Sajjadifar, Z. Arzehgar, A. Ghayuri, $Zn_3(BTC)_2$ as a Highly Efficient Reusable Catalyst for the Synthesis of 2-Aryl-1H-Benzimidazole, *Journal of the Chinese Chemical Society*, 65(2017) 205-211.
- [10] A. Sajjadi, R. Mohammadi, Fe_3O_4 magnetic nanoparticles (Fe_3O_4 MNPs): A magnetically reusable catalyst for synthesis of Benzimidazole compounds, *Journal of Medicinal and Chemical Sciences*, 2 (2019) 55-58.
- [11] M. Abdollahi-Alibeik, M. Moosavifard, $FeCl_3$ -Doped Polyaniline Nanoparticles as Reusable Heterogeneous Catalyst for the Synthesis of 2-Substituted Benzimidazoles, *Synthetic Communication*, 40 (2010) 2686-2695.
- [12] S. Ponnala, D. P. Sahu, Iodine-Mediated Synthesis of 2-Arylbenzoxazoles, 2-Arylbenzimidazoles, and 1,3,5-Trisubstituted Pyrazoles, *Synthetic Communication*, 36 (2006) 2189-2194.
- [13] F. H. Havaladar, G. Mule, B. Dabholkar, Novel Method for the Syntheses of 2-Substituted Benzimidazoles Using Ti(IV) Isopropoxide and Cumene Hydroperoxide, *Synthetic Communication*, 41 (2011) 2304-2308.
- [14] H. Sharghi, O. Asemani, R. Khalifeh, New One-Pot Procedure for the Synthesis of 2-Substituted Benzimidazoles, *Synthetic Communication*, 38 (2008) 1128-1136.
- [15] C. Mukhopadhyay, P. K. Tapaswi, PEG-mediated catalyst-free expeditious synthesis of 2-substituted benzimidazoles and bis-benzimidazoles under solvent-less conditions, *Tetrahedron Letters*, 49 (2008) 6237-6240.
- [16] J. Yuan, Z. Zhao, W. Zhu, H. Li, X. Qian, Y. Xu, New strategy for the synthesis of 2-phenylbenzimidazole derivatives with sodium perborate (SPB) as oxidant, *Tetrahedron*, 69 (2013) 7026-7030.
- [17] G. A. Molander, K. Ajayi, Oxidative Condensations To Form Benzimidazole-Substituted Potassium Organotrifluoroborates, *Organic Letters*, 14 (2012) 4242-4245.

- [18] S. Samanta, S. Das, P. Biswas, Photocatalysis by 3,6-Disubstituted-s-Tetrazine: Visible-Light Driven Metal-Free Green Synthesis of 2-Substituted Benzimidazole and Benzothiazole, *Journal of Organic Chemistry*, 78 (2013) 11184-11193.
- [19] S. M. Inamdar, V. K. More, S. K. Mandal, CuO nano-particles supported on silica, a new catalyst for facile synthesis of benzimidazoles, benzothiazoles and benzoxazoles, *Tetrahedron Letters*, 54 (2013) 579-583.
- [20] K. Bahrami, M. M. Khodaei, F. Naali, Mild and Highly Efficient Method for the Synthesis of 2-Arylbenzimidazoles and 2-Arylbenzothiazoles, *Journal of Organic Chemistry*, 73 (2008) 6835-6837.
- [21] B. Das, H. Holla, Y. Srinivas, Efficient (bromodimethyl)sulfonium bromide mediated synthesis of benzimidazoles, *Tetrahedron Letters*, 48 (2007) 61-64.
- [22] S. S. Panda, S. C. Jain, Synthesis of 2-Arylbenzimidazoles in Water, *Synthetic Communication*, 41 (2011) 729-735.
- [23] P. Gogoi, D. Konwar, An efficient and one-pot synthesis of imidazolines and benzimidazoles via anaerobic oxidation of carbon-nitrogen bonds in water, *Tetrahedron Letters*, 47 (2006) 79-82.
- [24] A. Dhakshinamoorthy, K. Kanagaraj, K. Pitchumani, Zn²⁺-K10-clay (clayzic) as an efficient water-tolerant, solid acid catalyst for the synthesis of benzimidazoles and quinoxalines at room temperature, *Tetrahedron Letters*, 52 (2011) 69-73.
- [25] Y. Riadi, R. Mamouni, R. Azzalou, M. E. Haddad, S. Routier, An efficient and reusable heterogeneous catalyst Animal Bone Meal for facile synthesis of benzimidazoles, benzoxazoles, and benzothiazoles, *Tetrahedron Letters*, 52 (2011) 3492-3495.
- [26] V. Kumar, D. G. Khandare, A. Chatterjee, M. Banerjee, DBSA mediated chemoselective synthesis of 2-substituted benzimidazoles in aqueous media, *Tetrahedron Letters*, 54 (2013) 5505-5509.
- [27] J. Lu, B. Yang, Y. Bai, Microwave Irradiation Synthesis of 2-Substituted Benzimidazole Using PPA as a Catalyst Under Solvent-Free Condition, *Synthetic Communication*, 32 (2002) 3703-3709.
- [28] S. Kaul, A. Kumar, B. Sain, A. K. Bhatnagar, Simple and Convenient One-Pot Synthesis of Benzimidazoles and Benzoxazoles using N,N-Dimethylchlorosulfitemethaniminium Chloride as Condensing Agent, *Synthetic Communication*, 37 (2007) 2457-2460.
- [29] H. Alinezhad, F. Salehian, P. Biparva, Synthesis of Benzimidazole Derivatives Using Heterogeneous ZnO Nanoparticles, *Synthetic Communication*, 42 (2012) 102-108.
- [30] P. C. Guillermo, B. A. Imelda, G. L. C. Jose, G. E. E. Jose, A. T. Cecilio, Synthesis of Benzimidazoles in Dry Medium, *Synthetic Communication*, 30 (2000) 2191-2195.
- [31] X. Wen, J. E. Bakali, R. Deprez-Poulain, B. Deprez, Efficient propylphosphonic anhydride (@T3P) mediated synthesis of benzothiazoles, benzoxazoles and benzimidazoles *Tetrahedron Lett*, 53 (2012) 2440-2443.
- [32] X. Jing, Q. Zhu, F. Xu, X. Ren, D. Li, C. Yan, Rapid one-pot preparation of 2-substituted benzimidazoles from esters using microwave conditions, *Synthetic Communication*, 36 (2006) 2597-2601.
- [33] L. Wang, S. Jia, H. Tian, C. Qian, An Efficient Procedure for the Synthesis of Benzimidazole Derivatives Using Yb(OTf)₃ as Catalyst Under Solvent-Free Conditions,

Synthetic Communication. 34(2004) 4265-4272.

[34] D. S. VanVliet, P. Gillespie, J. J Scicinski, Rapid one-pot preparation of 2-substituted benzimidazoles from 2-nitroanilines using microwave conditions, Tetrahedron Letters, 46 (2005) 6741-6743.

[35] G. Li, J. Wang, B. Yuan, D. Zhang, Z. Lin, P. Li, H. Huang, Dehydrogenative synthesis of benzimidazoles under mild conditions with

supported iridium catalysts, Tetrahedron Letter, 54 (2013) 6934-6936.

[36] T. B. Nguyen, L. Ermolenko, A. Al-Mourabit, Iron Sulfide Catalyzed Redox/Condensation Cascade Reaction between 2-Amino/Hydroxy Nitrobenzenes and Activated Methyl Groups: A Straightforward Atom Economical Approach to 2-Hetaryl-benzimidazoles and -benzoxazoles, Journal of the American Chemical Society, 135 (2013) 118-121.

HOW TO CITE THIS ARTICLE

Hridoydip Ranjan Dasgupta, Suvodip Mukherjee, Pranab Ghosh, One Pot Reductive Synthesis of Benzimidazole Derivatives from 2-Nitro Aniline and Aromatic Aldehydes Using Zn/NaHSO₃ in Water Medium, Prog. Chem. Biochem. Res., 4(1) (2021) 57-67.

DOI: [10.22034/pcbr.2020.245413.1126](https://doi.org/10.22034/pcbr.2020.245413.1126)

URL: http://www.pcbiochemres.com/article_118362.html

