

## Original Research Article

## Comparative Energy Dispersive X- Ray Fluorescence Analysis of *Mangifera Indica L.* Leaves in the Locality of Kachchh and Saurashtra

Foram H. Vaghela<sup>1\*</sup>, Tejal D. Bhatt<sup>1</sup>, Kanji D. Kachhot<sup>1</sup>, Chirag H. Dhamal<sup>1</sup>, Vijay R. Ram<sup>2</sup>, Hitendra S. Joshi<sup>1</sup>

<sup>1</sup>Department of Chemistry, Saurashtra University, Rajkot-360005, Gujarat, India

<sup>2</sup>Department of Chemistry, Kskv Kachchh University, Bhuj-370001, Gujarat India

## ARTICLE INFO

## Article history

Submitted: 2022-07-02

Revised: 2022-09-03

Accepted: 2022-09-25

Available online: 2022-09-29

Manuscript ID: PCBR-2207-1227

DOI: 10.22034/pcbr.2022.349997.1227

## KEYWORDS

*Mangifera Indica L.*

ED-XRF analysis

Kachchh Region

Saurashtra Region

Elements

## ABSTRACT

The aim of the present study was to determine the comparative investigation of chemical composition in the leaves of *Mangifera indica L.* in the region of Kachchh and Saurashtra. *Mangifera Indica L.* belongs to the anacardiaceae family. The leaves of *Mangifera Indica L.* were subjected to energy dispersive X-ray fluorescence (EDXRF) and analysed for different mineral compositions. As we know, XRF is one of the most sensitive, accurate, consistent, and non-destructive methods for analysing major and trace elements by using a single pressed pellet. During the analysis, it was found that in kachchh and Saurashtra region, *Mangifera Indica L.* leaves contain silicon, calcium, potassium, chlorine, sulphur, magnesium, aluminium, and iron were noted in higher amounts compared with that of other elements like strontium, manganese, titanium, bromine, zinc, barium, rubidium, nickel, silver, lead, molybdenum, etc. whereas the elements which were not detected in the leaves of *Mangifera Indica L.* are sodium, vanadium, uranium, mercury, silver, gold, etc. In comparison of Kachchh and Saurashtra region of *Mangifera Indica L.* leaves calcium, silicon, potassium, aluminium, phosphorus, and iron were high in the region of Saurashtra rather than the Kachchh leaves and other elements are presented in trace amount.

\* Corresponding author: Foram H. Vaghela

✉ E-mail: [drhsjoshichem@gmail.com](mailto:drhsjoshichem@gmail.com)

© 2022 by SPC (Sami Publishing Company)



GRAPHICAL ABSTRACT



1-Introduction

X-Ray fluorescence (XRF)[1][2] is an analytical method which is firstly use for the lead detection (Pb)[3] in paint in the 1970s for the abatement and exposure studies [4]. It has been since use for the environmental analysis like alloys [5], geological materials [6], glasses, and sediments [7] with very minimal sample preparation and treatment. During several times, XRF has been accepted by the environmental research community as a viable analytical tool through the efficiency of excitation of the radioisotope source [8] in conjunction with extremely sensitive

detectors [9] and other electronics. Therefore, it offers multi-element analytical capability [10], economy, high speed, and simplified operation where its advantages [11] and limitations [12] are well-understood.

*Mangifera Indica L.* (anacardiaceae) [13] is a large leafy tree valued mainly for its green and ripe fruit [14]. About 500 mango varieties have been reported in India [15]. It can grow up to 1530 meters high. The tree grows best in well-drained sandy loam; it does not grow well in heavy wet soil. The optimal soil pH should be between 5.2 and 7.5 [16].



Fig. 1. *Mangifera Indica L.* plant photograph



**Fig. 2.** Powder sample of *Mangifera Indica L.*, Kachchh Region



**Fig. 3.** Powder sample of *Mangifera Indica L.*, Saurashtra Region

Mango fruit has a high nutritional value and health benefits [17] due to important components [18]. Mango fruit contains phytochemicals [19] like polyphenol, pigments, phenolic, and volatile constituents [20]. They also contain carbohydrates and cellulose [21]. Mango fruit also give anti-oxidant activity [22].

In recent years, traditional medicines have been extracted from medicinal plants [23]. The use of mango as a traditional medicine to treat diabetes [24], inflammation [25], bacteria [26], fungi, and gastric protection [27] has been further studied.

### 1-1- Botanical classification [28]

- Kingdom: Plantae
- Subkingdom: Tracheobionta
- Superdivision: Spermatophyta
- Division: Magnoliophyta
- Class: Magnoliopsida
- Subclass: Rosidae
- Order: Sapindales
- Family: Anacardiaceae
- Genus: *Mangifera*
- Species: *M. indica*

### 1-2- Common names of different regions [28]

The common names of *Mangifera Indica L.* are as follow:

- Arabic : Mabaz
- Bengali : Am (Um)

- Chinese: Mi wang
- Danish: Mango, Mangofrugt, and Mangotrae
- Dutch: Manga, Mangga, Manja, and Mangoestanboom
- English: Mango
- French: Mangue and Manguier
- German: Indischer Mangobaum and Mango
- Greek: Magko and Mangko
- Gujarat: Ambo
- Hindi: Am, Ambi, and Amia
- Japanese: Anchar, Mangoo, and Mangou
- Kachchh: Ambo

## 2- Experimental

### 2-1- Materials and methods

#### 2-1-1- Sample preparation

Leaves of *Mangifera Indica L.* were collected in Kachchh and Saurashtra Regions of Gujarat in November December 2020. The leaves were sun dried to evaporate the water content from them. Then, it was grounded in mixture and with the help of pellet maker. Pellets from leaf samples were used for further elemental analysis in on-instrument energy dispersive X-ray fluorescence.

#### 2-1-2- Instrumental parameters

Rigaku elemental analyzer are benchtop X-ray fluorescence (XRF) with element range Na to U with Pd anode X-ray tube with a high

performance SDD detector by using NEX CG software.



**Figure 6.** Pellet of *Mangifera Indica L.*, Kachchh Region



**Figure 7.** Pellet of *Mangifera Indica L.*, Saurashtra Region

### 3- Results and Discussion

#### 3-1- Chemicals


The *Mangifera Indica L.* Leaves growing in the region between Northwest like India, Myanmar, and Bangladesh. *Mangifera Indica L.* was collected from Kachchh and Saurashtra districts and subjected to the X-ray fluorescence instrument for mineral analysis for the present study. Various and large amounts of mineral ions were found during the analysis.

The main constituent calcium (Ca) is a nutrient that all living organisms need, including humans.

It is the most abundant mineral in the body and is critical to bone health, found in the leaves of *M. Indica L.* in the Kachchh and Saurashtra Regions was 2.56% and 3.21%, respectively. Potassium (K) content, considered an important component for the body, was found in Kachchh and Saurashtra to be 1.07% and 1.20%, respectively. Silicon (Si), which is not only a good soil binder, but is also used for many industrial purposes, was also found in larger amounts in the leaves of *M. Indica L.* for Kachchh and Saurashtra at 3.13% and 2.14%, respectively found.

**Table 1.** The XRF analysis of *Mangifera Indica L.*, Kachchh Region


Sr. No.	Element	% Mass
1	Si	3.13
2	Ca	2.56
3	K	1.07
4	Cl	0.419
5	S	0.370
6	Mg	0.328
7	Al	0.308
8	P	0.203
9	Fe	0.0641
10	Sr	0.0233
11	Mn	0.0063
12	Ti	0.0042
13	Br	0.0031
14	Zn	0.0029
15	Cu	0.0018
16	Zr	0.0016
17	Ba	0.0008



**Fig. 4.** The XRF graph for mineral ion in the leaves of *Mangifera Indica L*, Kachchh Region

**Table 2.** The XRF analysis of *Mangifera Indica L*, Saurashtra Region

Sr. No.	Element	% Mass
1	Si	2.14
2	Ca	3.21
3	K	1.20
4	Cl	0.249
5	S	0.266
6	Mg	0.238
7	Al	0.312
8	P	0.228
9	Fe	0.112
10	Sr	0.0162
11	Mn	0.0058
12	Ti	0.0089
13	Br	0.0008
14	Zn	0.0029
15	Cu	0.0017
16	Zr	0.0012
17	Cr	0.0018
18	Rb	0.0007
19	Ni	0.0006
20	Ag	0.0001
21	Pb	0.0001
22	Mo	0.0001



**Fig. 5.** The XRF Graph for mineral ion in the leaves of *Mangifera Indica L.*, Saurashtra Region

The major component found in the leaves of *M. Indica L.* from Kachchh area is chlorine (0.419%), sulphur (0.370%), magnesium (0.328%), aluminium (0.308%), and phosphorus (0.203%) and for the Saurashtra area, chlorine (0.249%), sulphur (0.266%), magnesium (0.238%), aluminium (0.312%), phosphorus (0.228%), and Fe (0.112%) were noted.

Important findings from the leaves of *Mangifera Indica L.* were the presence of strontium (Sr), manganese (Mn), titanium (Ti), bromine (Br), zinc (Zn), copper (Cu), zirconium (Zr), barium (Ba), chromium (Cr), rubidium (Rb), nickel (Ni), silver (Ag), lead (Pb), and molybdenum (Mo) were found in trace amounts.

The elements which were undetected were Y, As, U, Na, Ru, Rh, Cd, Sn, Sb, Au, La, Ce, Pr, Nd, W, etc.

#### 4. Conclusion

The ED XRF method is a powerful technique for analysing various anions and cations found in the leaves of *Mangifera Indica L.* in the regions of Kachchh and Saurashtra in Gujarat, India. The components with higher values in Saurashtra are calcium, silicon, potassium, aluminium, phosphorus, and iron compared with the

Kachchh Region and other elements like chlorine, strontium, manganese, titanium, bromine, zinc, copper, zirconium, barium, chromium, rubidium, nickel, silver, lead, and molybdenum were found in traces.

#### Acknowledgments

The authors are thankful to the Department of Chemistry, Saurashtra University, Rajkot for providing the laboratory facilities and also thankful to Department Of Chemistry, Shree Shyamji Krishna Verma Kachchh University, Bhuj-370001.

#### Supporting Information

*Mangifera Indica L.* was found in Kachchh and Saurashtra Region of Gujarat. The XRF analysis data and experimental method are available in the supplementary file.

#### Compliance with Standards

This article does not contain any studies involving human participants performed by any authors and does not contain any studies involving animals performed by any of the author.

## Conflict of Interest

The authors report no conflicts of interest

## References

- [1] M.S. Shackley, An introduction to X-ray fluorescence (XRF) analysis in archaeology. X-ray fluorescence spectrometry (XRF) in geoarchaeology, New York, 3 (2011) 07-44.
- [2] T.D.T. Oyedotun, Evaluation of the effect of the fluxing agent over the purification of Columbite-Tantalite ore for the synthesis of Nb, Ta mixed oxides. *Geology, Ecology, and Landscapes*, 2 (2018) 148-154.
- [3] A. Al Maliki, A.K. Al-lami, H.M. Hussain, N. Al-Ansari, Comparison of portable X-ray fluorescence spectrometry and laboratory-based methods to assess the soil elemental composition: Applications for wetland soils. *Environmental Earth Sciences*, 76 (2017) 1-7.
- [4] V.G. Mihucz, et al., Removal of some elements from washed and cooked rice studied by inductively coupled plasma mass spectrometry and synchrotron based confocal micro-X-ray fluorescence. *Food Chemistry*, 121.1 (2010) 290-297.
- [5] Šatović, D., V. Desnica, and S. Fazinić. "Use of portable X-ray fluorescence instrument for bulk alloy analysis on low corroded indoor bronzes." *Spectrochimica Acta Part B: Atomic Spectroscopy* 89 (2013) 7-13.
- [6] Laubach, S. E., Lander, R. H., Criscenti, L. J., Anovitz, L. M., Urai, J. L., Pollyea, R. M., Pyrak-Nolte. "The role of chemistry in fracture pattern development and opportunities to advance interpretations of geological materials." *Reviews of Geophysics* 57.3 (2019)1065-1111.
- [7] Kenna, T. C., Nitsche, F. O., Herron, M. M., Mailloux, B. J., Peteet, D., Sritrairat, S., Baumgarten, J. Evaluation and calibration of a Field Portable X-Ray Fluorescence spectrometer for quantitative analysis of siliciclastic soils and sediments. *Journal of Analytical Atomic Spectrometry*, 26.2 (2011) 395-405.
- [8] Sychov, M., Kavetsky, A., Yakubova, G., Walter, G., Yousaf, S., Lin, Q., Bower, K. Alpha indirect conversion radioisotope power source. *Applied Radiation and Isotopes*, 66.2 (2008) 173-177.
- [9] Zuo, Y. Y., Ding, M., Bateni, A., Hoorfar, M., Neumann, A. W., Improvement of interfacial tension measurement using a captive bubble in conjunction with axisymmetric drop shape analysis (ADSA). *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 250.1-3 (2004) 233-246.
- [10] P.W. Abrahams, J.A. Entwistle, Robert A. Dodgshon. The Ben Lawers historic landscape project: simultaneous multi-element analysis of former settlement and arable soils by X-ray fluorescence spectrometry. *Journal of Archaeological Method and Theory*, 17.3 (2010) 231-248.
- [11] Carmona, N., Ortega-Feliu, I., Gomez-Tubio, B., Villegas, M. A. Advantages and disadvantages of PIXE/PIGE, XRF and EDX spectrometries applied to archaeometric characterisation of glasses. *Materials Characterization*, 61.2 (2010) 257-267.
- [12] I. Liritzis, and N. Zacharias. "Portable XRF of archaeological artifacts: current research, potentials and limitations." *X-ray fluorescence spectrometry (XRF) in geoarchaeology* (2011) 109-142.
- [13] Li, L., Wu, H. X., Ma, X. W., Xu, W. T., Liang, Q. Z., Zhan, R. L., Wang, S. B. Transcriptional mechanism of differential sugar accumulation in pulp of two contrasting mango (*Mangifera indica* L.) cultivars. *Genomics*, 112.6 (2020) 4505-4515.
- [14] Lauricella, M., Emanuele, S., Calvaruso, G., Giuliano, M., D'Anneo, A. Multifaceted health benefits of *Mangifera indica* L. (Mango): the inestimable value of orchards recently planted in Sicilian rural areas. *Nutrients*, 9.5 (2017) 525.

- [15] M. Srivastav, et al. New hyper-variable SSRs for diversity analysis in mango (*Mangifera indica* L.). *Indian Journal of Genetics and Plant Breeding*, 81.01 (2021) 119-126.
- [16] Cáceres-Mago, Karla, Alicia Cáceres, and Luis D. Llambí. Effects of nurse shrubs on symbioses between soil fungi and associated plants along a tropical alpine elevation gradient. *Alpine Botany*, (2021) 1-16.
- [17] Maldonado-Celis, M. E., Yahia, E. M., Bedoya, R., Landázuri, P., Loango, N., Aguillón, J., Guerrero Ospina, J. C. Chemical composition of mango (*Mangifera indica* L.) fruit: Nutritional and phytochemical compounds. *Frontiers in plant science*, 10 (2019) 1073.
- [18] R.N. Tharanathan, H.M. Yashoda, T.N. Prabha. Mango (*Mangifera indica* L.), The king of fruits"—An overview. *Food Reviews International*, 22.2 (2006) 95-123.
- [19] Kumar, M., Saurabh, V., Tomar, M., Hasan, M., Changan, S., Sasi, M. Mekhemar, M. Mango (*Mangifera indica* L.) leaves: Nutritional composition, phytochemical profile, and health-promoting bioactivities. *Antioxidants*, 10.2 (2021): 299.
- [20] Gupta, A. K., Gurjar, P. S., Beer, K., Pongener, A., Ravi, S. C., Singh, S., Verma, D. K. A review on valorization of different byproducts of mango (*Mangifera indica* L.) for functional food and human health. *Food Bioscience*. (2022): 101783.
- [21] Cárdenas-Pérez, S, Chanona-Pérez, J. J., Güemes-Vera, N., Cybulska, J., Szymanska-Chargot, M., Chylinska, M Zdunek, A. Structural, mechanical and enzymatic study of pectin and cellulose during mango ripening. *Carbohydrate polymers*. 196 (2018) 313-321.
- [22] A. Bharathi, K. Prabhu, S. Rajan. Bactericidal effect of tender and matured seed kernel of *Mangifera indica* a comparative study. *Int J Life Sci Res.*, 7 (2019) 175-80.
- [23] R.A. Dar, M. Shahnawaz, P.H. Qazi. General overview of medicinal plants: A review. *The Journal of Phytopharmacology*, 6.6 (2017) 349-351.
- [24] Visentin, A. P. V., Colombo, R., Scotton, E., Fracasso, D. S., da Rosa, A. R., Branco, C. S., Salvador, M. Targeting inflammatory-mitochondrial response in major depression: current evidence and further challenges. *Oxidative Medicine and Cellular Longevity*, 2020 (2020) 2972968.
- [25] Kim, H., Castellon-Chicas, M. J., Arbizu, S., Talcott, S. T., Drury, N. L., Smith, S., Mertens-Talcott, S. U. Mango (*Mangifera indica* L.) polyphenols: Anti-inflammatory intestinal microbial health benefits, and associated mechanisms of actions. *Molecules*, 26.9 (2021) 2732.
- [26] N.A. Chugh, S. Bali, A. Koul. Integration of botanicals in contemporary medicine: road blocks, checkpoints and go-ahead signals. *Integrative medicine research*, 7.2 (2018) 109-125.
- [27] Ponpandian, S. Vinoth, and A. Egbert Selwin Rose. Arbuscular Mycorrhizal Fungi Associated With Some Agroforestry Tree Species From Eastern Ghats of Tamil Nadu. *Indian Journal of Scientific Research*. (2018) 6-12.
- [28] G.M. Masud Parvez. Current advances in pharmacological activity and toxic effects of various *Capsicum* species. *Int. J. Pharm. Sci. Res.*, 8.5 (2017) 1900-1912.

#### HOW TO CITE THIS ARTICLE

Foram H. Vaghela, Tejal D. Bhatt, Kanji D. Kachhot, Chirag H. Dhamal, Vijay R. Ram, Hitendra S. Joshi Comparative Energy Dispersive X- Ray Fluorescence Analysis of *Mangifera Indica* L. Leaves In the Locality of Kachchh and Saurashtra. *Prog. Chem. Biochem. Res*, 5(3) (2022)254-261

DOI: 10.22034/pcbr.2022.349997.1227

URL: [http://www.pcbiochemres.com/article\\_158126.html](http://www.pcbiochemres.com/article_158126.html)

